

Toppan Best-set Premedia Limited	
Journal Code: JZO	Proofreader: Mony
Article No: JZO12018	Delivery date: 30 Jan 2013
Page Extent: 11	

Activity patterns and temporal avoidance by prey in response to Sunda clouded leopard predation risk

J. Ross^{1*,2}, A. J. Hearn^{1*,2}, P. J. Johnson¹ & D. W. Macdonald¹

¹ Wildlife Conservation Research Unit (WildCRU), Department of Zoology, University of Oxford, Oxford, UK

² Global Canopy Programme, Oxford, UK

Keywords

activity patterns; circular statistics; overlap coefficient; Sunda clouded leopard; ungulate.

Correspondence

Joanna Ross, Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, The Recanati-Kaplan Centre, Tubney House, Abingdon Road, Tubney, Abingdon OX13 5QL, UK.
Email: joanna.ross@zoo.ox.ac.uk

Editor: Nigel Bennett

*Current address

Received 6 August 2012; revised 30 December 2012; accepted 16 January 2013

doi:10.1111/jzo.12018

Introduction

Temporal interactions between predators and prey can be complex. Predators are expected to maximize their hunting efficiency (their intake per unit effort) by exploiting periods of prey vulnerability, while at the same time limiting competition with other predators. For example, carnivores specialized in burrowing such as badgers *Meles meles* dig prey from their underground refuges, the same prey species being hunted above ground by predators relying on speed (red fox, *Vulpes vulpes*) or stealth (European lynx, *Lynx pardinus*) (Fedriani, Palomares & Delibes, 1999). Prey species have to trade optimal feeding opportunities, finding mates and reducing competition with conspecifics against avoiding encounters with predators. Prey may avoid predators by temporal or spatial shifts. Valeix *et al.* (2009) describe how African ungulates altered their space use in response to the risk of lion predation; likewise, Fedriani *et al.* (1999) report that the red fox avoids conflict with the larger European lynx by a differential use of space. Bowyer *et al.* (1999) describe how female ungulates move to areas of lower predation risk before giving birth. Avoidance by shifting activity patterns is described, for example, by Eccard *et al.* (2008) and Gliwicz & Drabowski (2008) who found that bank voles *My. glareolus* and root

Abstract

Little is known about the activity patterns of Bornean ungulates, or the temporal interactions of these species with the Sunda clouded leopard *Neofelis diardi*. In this study, we use photographic capture data to quantify the activity patterns for the Sunda clouded leopard and six potential prey species: bearded pig *Sus barbatus*, Bornean yellow muntjac *Muntiacus atherodes*, red muntjac *Muntiacus muntjak*, lesser mouse deer *Tragulus kanchil*, greater mouse deer *Tragulus napu*, and sambar deer *Rusa unicolor*, and to calculate the overlap in activity patterns between these species. This is the first insight into the temporal interactions between the Sunda clouded leopard and its potential prey. Sunda clouded leopards' activity patterns overlapped most with those of sambar deer and greater mouse deer. In the absence of clouded leopards, we report a significant difference in activity patterns for bearded pigs which show greater nocturnal activity in the absence of this predator. This suggests that bearded pigs may be prey species for clouded leopards and they are capable of altering their activities in response to this risk.

voles *Mi. oeconomus*, respectively, became nocturnal in the presence of the diurnal least weasel (*Mu. nivalis*). A similar shift was described by Fenn & Macdonald (1995) who reported that a population of Norway rats *Rattus norvegicus* was diurnal (rather than the typical nocturnal) when facing a high risk of nocturnal predation by red foxes.

However, as Valeix *et al.* (2009) observed, prey species' behavioural plasticity varies and therefore, conversely, and provided the effect of competition is not dominant, we might expect predator activity to be often in phase with the periods when those potential prey species are most vulnerable to their method of predation. For some predators, this may result in their activity patterns mirroring those of their prey; while we would not expect this to be a general phenomenon, it may be commonest for felids that hunt principally using visual cues (Sunquist & Sunquist, 2002). Harmsen *et al.* (2011), for example, have reported this mirroring effect of predator and prey for both jaguar *Pa. onca* and puma *Pu. concolor* and their respective main prey species in a rainforest environment. Jenny & Zuberbuhler (2005) report that leopard *Panthera pardus* activity patterns, although varying across Africa, remain in phase with its main prey. Likewise, Kawanishi & Sunquist (2004) found that tigers in Taman Negara National Park, Malaysia, were mainly diurnal, an activity pattern that

1 corresponds with that of their most important prey species.
2 There are some conspicuous counter-examples however, for
3 example, the lion *Panthera leo* is a primarily nocturnal hunter
4 but is principally dependent on plains zebra *Equus quagga* and
5 common wildebeest *Connochaetes taurinus* as prey (Mills &
6 Shenk, 1992), species that are active diurnally.

7 Like all felids, the Sunda clouded leopard *Neofelis diardi*
8 (hereafter clouded leopard) is an obligate carnivore, and is
9 predominantly nocturnal, with some crepuscular activity
10 (Cheyne & Macdonald, 2011; Hearn *et al.*, submitted). In the
11 absence of any larger carnivore in Borneo, this time tabling
12 may function to maximize hunting opportunities rather than
13 as a tactic to avoid interspecific conflicts.

14 While detailed information is currently lacking, the
15 clouded leopard's diet is known to include sambar deer *Rusa*
16 *unicolor*, muntjacs *Muntiacus* spp., bearded pig *Sus barbatus*,
17 mouse deer *Tragulus* spp. (Rabinowitz, Andau & Chai, 1987;
18 Mohamed, Samejima & Wilting, 2009), porcupines (Gordon
19 & Stewart, 2007) and primates (Yeager, 1991; Matsuda,
20 Tuuga & Higashi, 2008). The patterns of activity for most of
21 these ungulate prey species in Borneo remain unclear as do
22 the temporal interactions with clouded leopards. Elsewhere,
23 the ungulate guild composition differs as does predation
24 pressure, potentially shaping different patterns of activity for
25 these ungulates. Across their ranges, the sambar deer is noc-
26 turnal (Schaller, 1967; Payne & Francis, 1998; Kawanishi
27 & Sunquist, 2004), the red muntjac *Muntiacus muntjak* is
28 diurnal (Payne & Francis, 1998; Kawanishi & Sunquist,
29 2004) or nocturnal (Kamler *et al.*, 2012), the yellow
30 muntjac *Muntiacus atherodes* is diurnal (Payne & Francis,
31 1998), the lesser mouse deer *Tragulus kanchil* is crepuscular
32 (Matsubayashi, Bosi & Kohshima, 2003) or arrhythmic
33 (Payne & Francis, 1998), and the bearded pig mostly diurnal
34 (Payne & Francis, 1998). In this study, we use photographic
35 capture data from Borneo to quantify the activity patterns
36 for six ungulate species: bearded pig *S. barbatus*, Bornean
37 yellow muntjac *M. atherodes*, red muntjac *M. muntjak*, lesser
38 mouse deer *T. kanchil*, greater mouse deer *Tragulus napu*,
39 and sambar deer *R. unicolor*, and investigate the temporal
40 segregation between them. Temporal partitioning among
41 sympatric species of Muridae has been reported by Meek,
42 Zewe & Falzon (2012) who speculate that this may reduce
43 competition for food resources, but this has not previously
44 been investigated for an ungulate guild. We compare activity
45 cycles between males and females for clouded leopards and
46 also for bearded pigs. For the other ungulate species, this
47 subdivision was not possible. Either we could not reliably
48 distinguish males from females in the photographs while
49 retaining sufficient sample sizes, or in the case of the yellow
50 muntjac, males and females do not travel independently of
51 one another. We then compare the activity patterns of Sunda
52 clouded leopard with those of the sympatric ungulates to
53 draw inferences about principal prey species. Finally, we
54 compare ungulate activity cycles in forests where clouded
55 leopards were detected with those in which this predator
56 appears to be absent. Clouded leopards may exert a top-
57 down effect on the activity cycles of ungulates. In the
58 absence of this apex predator, some species may be released

59 from a predominantly diurnal activity cycle with conse-
60 quences for the balance of competition with other ungulates.

61 We hypothesize firstly (Hypothesis 1) that if ungulates are
62 vulnerable to predation by clouded leopard they will be more
63 diurnal in areas where clouded leopards are present, compared
64 with areas where clouded leopards are not present. If ungulate
65 activity is not affected by the presence of clouded leopard, one
66 plausible explanation is that the prey are not vulnerable
67 because, for example, they are too small or too large to be
68 hunted efficiently. Secondly (Hypothesis 2), we hypothesize
69 that contest competition among ungulates will lead to lower
70 temporal overlap for those species that exploit similar food
71 resources that are patchily distributed, such as fruit.

72 **Methods**

73 **Study sites**

74 As part of an ongoing assessment of Sunda clouded leopard
75 abundance and responses to habitat modification, six forest
76 areas (Danum Valley Conservation Area, Ulu Segama, Malua
77 and Kabili-Sepilok Forest Reserves, Tabin Wildlife Reserve,
78 and the Lower Kinabatangan Wildlife Sanctuary) in Sabah,
79 Malaysian Borneo, were surveyed with camera traps between
80 November 2006 and August 2011. Together, these areas cover
81 primary and selectively logged lowland Dipterocarp rainforest
82 and selectively logged predominantly riverine rainforest.
83 Clouded leopards were detected in all survey areas except
84 Kabili-Sepilok Forest Reserve (hereafter Sepilok), presenting
85 the opportunity to investigate ungulate activity patterns in the
86 absence of a natural predator.

87 **Field methods**

88 We established 363 camera stations across the study sites. We
89 set up cameras on old logging roads and both animal and
90 man-made trails; they were positioned *c.* 40–50 cm above the
91 ground with locations separated by *c.* 1.5–2.0 km. In all areas,
92 we used passive infrared digital camera traps of varying
93 models: Snapshot Sniper P41 (Snapshot Sniper LLC, OK,
94 USA), Cuddeback Capture (Non Typical Inc., WI, USA),
95 Bushnell Trophycam 2010 (Bushnell Corporation, KS, USA),
96 Reconyx HC500 (Reconyx Inc., WI, USA) and Panthera V3
97 (Panthera, New York, NY, USA). At all locations, we set the
98 cameras to operate continuously for 24 h of each day with
99 either a white Xenon flash (Snapshot Sniper, Cuddeback
100 Capture and Panthera V3) or infrared LED flash (Bushnell
101 Trophycam 2010 and Reconyx HC500) firing under low light
102 levels. All models of camera recorded the time and date of
103 each photographic event.

104 **Data analysis**

105 Following van Schaik & Griffiths (1996), we assumed that
106 when an animal of a given species passed in front of the
107 camera, the likelihood of obtaining a photo-capture was equal
108 throughout both night and day. Additionally, we assumed
109
110
111
112

Table 1 The number of independent records of Sunda clouded leopard *Neofelis diardi* and six presumed prey species from each survey area. The number of trap days in each area is indicated in parentheses. The area covered by camera traps in each site is indicated as is the sample size for males and females where relevant

Species	Survey area					
	Danum Valley (6228)	Ulu Segama (4154)	Malua (3343)	Tabin (6172)	Kinabatangan (3997)	Sepilok (3755)
Area of camera grid (km ²)	66.26	60.07	102.80	160.28	111.0	52.4
Bearded pig <i>Sus barbatus</i>	379 (153 male, 36 female)	619 (210 male, 185 female)	828 (215 male, 57 female)	1056 (202 male, 199 female)	1308 (120 male, 275 female)	167 (41 male, 19 female)
Bornean yellow muntjac <i>Muntiacus atherodes</i>	542	126	134	1189	6	112
Red muntjac <i>Muntiacus muntjak</i>	100	167	24	70	0	0
Lesser mouse deer <i>Tragulus kanchil</i>	526	43	207	952	32	340
Greater mouse deer <i>Tragulus napu</i>	308	162	202	173	17	250
Sambar deer <i>Rusa unicolor</i>	291	214	608	587	90	21
Sunda clouded leopard	22 (11 male, 10 female)	104 (87 male, 16 female)	10 (8 male, 2 female)	41 (36 male, 4 female)	14 (7 male, 6 female)	0

that the manner in which clouded leopards hunt would not affect detections. The date and time of each event were extracted from the EXIF data of each photograph. To reduce bias caused by repeat detections of the same animal, one record of each species per hour per camera site was considered as a detection event for each 24-h period, and subsequent records were eliminated.

The timing of sunrise and sunset in Sabah is fairly consistent throughout the year and across study sites. Sunrise is roughly 06:00 h and sunset 18:00 h, local time (GMT + 8). We categorized nocturnal activity as that which fell between 19:00 and 05:00 h and diurnal as between 07:00 and 17:00 h; the remaining periods we classed as crepuscular.

Statistical modelling of activity patterns was effected using circular techniques (Zar, 1996). We constructed models predicting daily activity as a function of continuous trigonometric predictor variables describing one and two complete cycles in a 24-h period ($\sin\theta$, $\cos\theta$, and $\sin 2\theta$, $\cos 2\theta$ with $\theta = \pi t/24$, where t is time in hours). For sympatric species, we first tested whether activity cycles differed among species using a categorical predictor with levels indicating species. For models testing for an effect of the presence of clouded leopard, we used a categorical term indicating site (firstly using six levels, one of which, Sepilok, was thought to not have clouded leopards). To investigate this further, we compared all sites with known presence of clouded leopard using five levels and, finally, we pooled the data from these five sites to compare with Sepilok using two levels. Patterns were compared between the sexes for bearded pigs and clouded leopards using a similar method. In the presence of interaction (significance of trigonometric \times categorical terms) indicating evidence for varying periodicity, we fitted separate models for each level of the category (e.g. species). These models identified the amplitude and number of cycles within each 24-h period (see de Bruyn *et al.*, 2003). Patterns were visualized using the package *Plotrix* in R (Lemon, 2006).

We also calculated a coefficient of overlap between the activity patterns of each ungulate and Sunda clouded leopard from pooled forest data, including a novel bootstrap procedure, to estimate the precision of this measure following Ridout & Linkie (2009) and Linkie & Ridout (2011).

Results

We accumulated a total of 27 649 trap days across all study areas. The numbers of independent photo-captures for each species (Table 1) were considerably higher than previously reported for similar studies (e.g. Linkie & Ridout, 2011).

Overall activity patterns

There was strong evidence that activity patterns differed among species ($\cos x_1 * \text{species}$: $F_{5, 114} = 25.7$, $P < 0.001$; $\sin x_1 * \text{species}$: $F_{5, 114} = 2.6$, $P = 0.03$). We therefore examined the cycles separately for each species (Fig. 1). Sunda clouded leopards showed a strongly nocturnal pattern of activity. Greater mouse deer and sambar deer were also clearly nocturnal. In contrast, both yellow and red muntjacs were diurnal, although red muntjac showed a higher peak in activity at dawn than did yellow muntjac. The lesser mouse deer was largely crepuscular and the bearded pig showed no distinct overall pattern. More detailed analyses of bearded pig activity, however, revealed patterns with clear cycles related to sex (Fig. 2).

Sex differences

Bearded pigs exhibited clear intersexual differences in activity ($\cos x_1 * \text{sex}$: $F_{1, 38} = 11.1$, $P = 0.002$) and there was weaker evidence for a difference between male and female clouded leopards ($\cos x_2 * \text{sex}$: $F_{1, 38} = 4.0$, $P = 0.054$). Male bearded pigs exhibited a bimodal, crepuscular activity cycle, whereas




Figure 1 Periodic regression analysis of clouded leopard and ungulate activity patterns. Actual data are shown as bars and the model is plotted as a line. The plot is divided into hours with the response axis showing the percentage of detections in each hour. (a) Sunda clouded leopard *Neofelis diardi*, (b) bearded pig *Sus barbatus*, (c) sambar deer *Rusa unicolor*, (d) red muntjac *Muntiacus muntjak*, (e) Bornean yellow muntjac *Muntiacus atherodes*, (f) lesser mouse deer *Tragulus kanchil*, (g) greater mouse deer *Tragulus napu*.

females were predominantly diurnal, with their activity peaking around midday. Both male and female clouded leopards showed predominantly nocturnal cycles, but the males were proportionally more active than females at dusk and dawn (Fig. 2).

Absence of clouded leopards

In Sepilok Forest Reserve, where clouded leopards were likely to have been absent, the activity cycles of bearded pigs differed from those in forests where clouded leopards were present, most markedly for male bearded pigs, to a lesser extent female bearded pigs and also greater mouse deer. Both male and female pigs showed a higher level of nocturnality in this area compared with the forests in which clouded leopards were present. Greater mouse deer were primarily nocturnal in both

areas, but showed proportionally more activities at dusk and in the earlier part of the night in the absence of clouded leopards (Fig. 3). In contrast, there was little evidence that the other ungulate species were affected by the absence of clouded leopard (Fig. 3, Table 2). The comparisons between the remaining five sites where clouded leopards were present did not show any significant differences in activity patterns for yellow muntjac. There were significant differences for male and female muntjac. There were significant differences for male and female bearded pigs, greater mouse deer and sambar deer (Table 3). In the presence of clouded leopards, the activity cycle of male pigs was either crepuscular or strongly diurnal and female pigs were predominantly diurnal, although in Malua the peak of activity was in the late afternoon and very early evening. Greater mouse deer maintained a predominantly nocturnal activity cycle across study areas, but showed a strongly bimodal pattern with peaks at dawn and dusk in

J. Ross *et al.*

Activity of Sunda clouded leopards and their prey




Figure 1 Continued.




Figure 2 Differences between males (solid line) and females (dashed line) for (a) bearded pigs *Sus barbatus* and (b) Sunda clouded leopards *Neofelis diardi*. The plot is divided into hours with the response axis showing the percentage of detections in each hour.

Activity of Sunda clouded leopards and their prey




Figure 3 Differences in areas with clouded leopards (solid line) and without (dashed line). (a) Male pigs, (b) female pigs, (c) greater mouse deer *Tragulus napu*, (d) lesser mouse deer *Tragulus kanchil*, (e) Bornean yellow muntjac *Muntiacus atherodes*, (f) sambar deer *Rusa unicolor*. The plot is divided into hours with the response axis showing the percentage of detections in each hour.

Table 2 Differences in ungulate activity patterns between areas with (pooled data) and without clouded leopards

Model	d.f.	Sum square	Mean square	F	P
Male bearded pigs					
sinx1*clouded leopard	1	2.606	2.606	0.5370	0.4681670
sinx2*clouded leopard	1	24.082	24.082	4.9621	0.0319063 ^a
cosx1*clouded leopard	1	91.464	91.464	18.8466	0.0001013 ^b
cosx2*clouded leopard	1	2.143	2.143	0.4416	0.5103666
Residuals	38	184.417	4.853		
Female bearded pigs					
sinx1*clouded leopard	1	0.02	0.022	0.0017	0.967210
sinx2*clouded leopard	1	41.75	41.746	3.2778	0.078136
cosx1*clouded leopard	1	6.12	6.120	0.4805	0.492392
cosx2*clouded leopard	1	12.85	12.849	1.0089	0.321530
Residuals	38	483.96	12.736		
Greater mouse deer <i>Tragulus napu</i>					
sinx1*clouded leopard	1	12.340	12.340	3.3130	0.07662
sinx2*clouded leopard	1	12.621	12.621	3.3887	0.07346
cosx1*clouded leopard	1	1.597	1.597	0.4287	0.51657
cosx2*clouded leopard	1	3.651	3.651	0.9804	0.32837
Residuals	38	141.533	3.725		
Lesser mouse deer <i>Tragulus kanchil</i>					
sinx1*clouded leopard	1	10.84	10.838	0.7786	0.383116
sinx2*clouded leopard	1	0.20	0.197	0.0142	0.905832
cosx1*clouded leopard	1	21.24	21.239	1.5259	0.224315
cosx2*clouded leopard	1	0.64	0.641	0.0460	0.831292
Residuals	38	528.93	13.919		
Bornean yellow muntjac <i>Muntiacus atherodes</i>					
sinx1*clouded leopard	1	0.16	0.16	0.0207	0.886334
sinx2*clouded leopard	1	0.09	0.09	0.0119	0.913662
cosx1*clouded leopard	1	0.14	0.14	0.0188	0.533400
cosx2*clouded leopard	1	3.03	3.03	0.3951	0.891590
Residuals	38	291.64	7.67		
Sambar deer <i>Rusa unicolor</i>					
sinx1*clouded leopard	1	0.35	0.346	0.0272	0.86985
sinx2*clouded leopard	1	6.00	5.995	0.4717	0.49637
cosx1*clouded leopard	1	33.39	33.393	0.0935	0.76139
cosx2*clouded leopard	1	1.19	1.189	2.6274	0.11330
Residuals	38	482.96	12.710		

^a••.
^b••.

Tabin. Similarly, sambar deer are nocturnal in all areas, but their activity peaks at dawn in Malua.

Activity overlap

The highest level of temporal overlap between ungulates (0.91) was observed for the greater mouse deer and sambar deer, which both showed a strongly nocturnal activity pattern (Fig. 4). The overlap between yellow and red muntjacs was also high (0.87). The lowest levels of overlap were between greater mouse deer and red muntjacs (0.32) and greater mouse deer and yellow muntjacs (0.27).

Both male and female clouded leopards overlapped most strongly with greater mouse deer, with overlap coefficients of 0.70 and 0.85, respectively (Fig. 4a and b). There was also high

overlap between both male and female clouded leopards and sambar deer (0.67 with male clouded leopards and 0.79 with females). Conversely, there was low overlap between the predominantly nocturnal clouded leopards and the diurnal muntjacs (0.24 and 0.19 with yellow muntjac, and 0.29 and 0.23 with red muntjac, for male and female clouded leopards, respectively). Overlap was also low between male and female clouded leopards and the crepuscular lesser mouse deer, with coefficients of 0.48 and 0.41, respectively. The overlap coefficient was lower for female clouded leopards and female bearded pigs than male pigs (0.40 and 0.47, respectively). Male clouded leopards overlapped more strongly with both male and female bearded pigs (0.64 and 0.52, respectively).

While the only species with statistically significantly different activity cycles between Sepilok and the other forest areas was bearded pig, and to a lesser extent greater mouse deer, the

Table 3 Differences in ungulate activity patterns between all areas where clouded leopards are present

Model	d.f.	Sum square	Mean square	F	P
Male bearded pigs					
sinx1*site	4	23.65	5.912	1.2706	0.2870099
sinx2*site	4	10.77	2.693	0.5788	0.6786860
cosx1*site	4	102.43	25.607	5.5036	0.0004986
cosx2*site	4	16.07	4.018	0.8636	0.4888376
Residuals	95	442.02	4.653		
Female bearded pigs					
sinx1*site	4	66.77	16.694	2.6973	0.035290
sinx2*site	4	10.12	2.531	0.4089	0.801805
cosx1*site	4	32.92	8.229	1.3296	0.264612
cosx2*site	4	34.48	8.620	1.3928	0.242334
Residuals	95	587.96	6.189		
Greater mouse deer <i>Tragulus napu</i>					
sinx1*site	4	51.92	12.98	1.2377	0.30021
sinx2*site	4	20.53	5.13	0.4893	0.74353
cosx1*site	4	124.14	31.03	2.9593	0.02368
cosx2*site	4	44.24	11.06	1.0546	0.38344
Residuals	95	996.27	10.49		
Lesser mouse deer <i>Tragulus kanchil</i>					
sinx1*site	4	33.25	8.31	0.5151	0.72482
sinx2*site	4	27.62	6.91	0.4279	0.78814
cosx1*site	4	138.06	34.52	2.1387	0.08196
cosx2*site	4	61.19	15.30	0.9480	0.43990
Residuals	95	1533.17	16.14		
Sambar deer <i>Rusa unicolor</i>					
sinx1*site	4	4.94	1.24	0.3231	0.861858
sinx2*site	4	4.14	1.03	0.2704	0.96358
cosx1*site	4	50.39	12.60	3.2929	0.014222
cosx2*site	4	23.57	5.89	1.5399	0.196885
Residuals	95	363.44	3.83		
Bornean yellow muntjac <i>Muntiacus atherodes</i>					
sinx1*site	4	66.55	16.64	0.7834	0.5387561
sinx2*site	4	28.92	7.23	0.3404	0.8500837
cosx1*site	4	13.82	3.46	0.1627	0.9567116
cosx2*site	4	30.16	7.54	0.3550	0.8399814
Residuals	95	2017.70	21.24		
Red muntjac <i>Muntiacus muntjak</i>					
sinx1*site	4	25.93	6.48	0.6163	0.606526
sinx2*site	4	24.50	6.13	0.5825	0.628286
cosx1*site	4	7.24	1.81	0.1721	0.914931
cosx2*site	4	42.90	10.73	1.0200	0.388629
Residuals	95	1065.62	11.22		

collective shifts in activity patterns resulted in some clearer differences in overlaps among the ungulates, there are numerous pairs of taxa where the confidence intervals for overlaps with and without clouded leopard do not overlap (Fig. 5). Sambar deer, for example, overlapped more with all other species apart from greater mouse deer in the absence of clouded leopards. Other clear effects included that in the absence of clouded leopards male and female pig activity and yellow muntjac overlapped less, and that conversely there were more overlaps between male pigs and greater mouse deer, and between greater mouse deer and yellow muntjac.

Discussion

This is one of the first studies to quantify the activity patterns of Bornean tropical forest ungulates and is the first to measure the overlap between the activity cycles of the Sunda clouded leopard and some of its potential prey species. We show a significant shift in bearded pig activity patterns in the absence of clouded leopards, suggesting that these activity cycles may be shaped, at least in part, by the risk of predation by clouded leopards. We also reveal a high temporal overlap between the clouded leopard and both sambar deer and greater mouse deer.

Activity patterns of Bornean ungulates and overlap with clouded leopards

We found bearded pigs to be predominantly diurnal (females) or crepuscular (males). A similar species, *Sus scrofa*, was also found to be diurnal in Sumatra (Linkie & Ridout, 2011). We found a significant shift in activity patterns for male pigs between areas of differing predation pressure. Male pigs were almost exclusively nocturnal where clouded leopards were

absent. It may be more cost effective for pigs, in terms of energy expenditure, to forage and move at night when temperatures are lower, but the risk of predation may be a stronger driving force and prevent this in areas where clouded leopards are present. This driving force may be stronger for female than for male pigs. Female bearded pigs showed a higher proportion of diurnal activities than did male pigs and showed a weaker shift in cycle in the absence of clouded leopards. Most pigs identified as females in this study were recorded with young, or were lactating; diurnality may be a strategy to avoid encountering clouded leopards during the period when piglets are most vulnerable to predation from this felid.

While there may be other contributory factors shaping the pigs' activity patterns that we are not accounting for, this activity shift for bearded pigs suggests that they may be prey species for clouded leopards, as it was only in Sepilok where the pigs were active nocturnally. The apparent behavioural plasticity of the bearded pig, allowing increased diurnality in the presence of clouded leopards, may be a tactic to avoid predation. Similarity in activity patterns between predator and prey is, by itself, not compelling evidence that a predator relies on any prey species. Felids such as jaguars and pumas may have activity patterns in phase with their main prey (Harmsen *et al.*, 2011) or, as with lions, have cycles that oppose those of their prey (Mills & Shenk, 1992). Predator-prey size relationships are also known to be important in shaping trophic links (Owen-Smith & Mills, 2008). Bearded pigs fall with the mass range that clouded leopards are likely to take based on prey mass to predator mass allometries (e.g. Carbone *et al.*, 1999; Macdonald, Mosser & Gittleman, 2010). Although adult pigs are at the greater end of this mass range, clouded leopards might target juvenile animals.

The weak evidence for activity shifts for the other ungulates between areas of different predation threat raises questions about their capacity to respond to contours in the landscape of fear. In Borneo, opportunistic observations of clouded leop-




Figure 4 The overlap between each ungulate species and (a) male clouded leopards and (b) female clouded leopards. Error bars show 95% bootstrap confidence intervals.




Figure 5 The difference in overlap between each pair of ungulates in areas with clouded leopards (grey bars) and without (white bars). Error bars show 95% bootstrap confidence intervals. BP = bearded pig *Sus barbatus*, BYM = Bornean yellow muntjac *Muntiacus atherodes*, LMD = lesser mouse deer *Tragulus kanchil*, GMD = greater mouse deer *Tragulus napu*. The dyads are ordered with greater shifts in overlap to the right.

ards with kills (Rabinowitz *et al.*, 1987) or the discovery of dead animals, presumed to have been killed by clouded leopards (Mohamed *et al.*, 2009), indicate that sambar deer, mouse deer species and muntjac species are also preyed on by clouded leopards, although we do not know their relative importance in its diet. Threat of predation by clouded leopards may be a weaker force in shaping their activity patterns than that of other predators, or avoiding contest competition for instance. Yellow and red muntjacs were diurnal as both Payne & Francis (1998) and Kawanishi & Sunquist (2004) have observed. For the yellow muntjac, this pattern held regardless of predation pressures (we could not assess this effect for the red muntjac due to small sample sizes). This presumed adaptation for diurnality may prevent a shift in the activity of yellow muntjacs, perhaps due to poor vision at night. A primarily nocturnal activity pattern for muntjacs in Lao People's Democratic Republic was, however, described by Kamler *et al.* (2012) who speculated that this was due to predation by the diurnal dhole *Cuon alpinus*. We also found high overlap between the two muntjacs' activity cycles. Both muntjacs have largely frugivorous diets, but red muntjacs are thought to browse more than do yellow (Barrette, 1977), which may explain the lack of temporal segregation. The low overlap between the muntjacs and the greater mouse deer may also be partly explained by diet; the greater mouse deer also eats fruits (Payne & Francis, 1998) and so this may be a strategy on the part of the mouse deer to avoid contest competition with the larger species. We describe a crepuscular activity cycle for the lesser mouse deer, a pattern also found by Matsubayashi *et al.* (2003), and this was not affected by the presence of clouded leopards. It is likely that to avoid contest competition with the larger nocturnal greater mouse deer, the activity of the lesser mouse deer, which is also frugivorous (Payne & Francis, 1998), is restricted to a crepuscular cycle. Consequently, temporal overlap between the two mouse deer species was relatively low. We found sambar deer to be strongly nocturnal corroborating Schaller (1967) and Kawanishi & Sunquist (2004). In this latter study, the activity pattern of the sambar deer may have been shaped by avoidance of the diurnal tiger. The highest coefficient of activity overlap was found between the sambar deer and the greater mouse deer. These species have dissimilar diets; the sambar deer is a mixed feeder, but predominantly a grazer and browser (Payne & Francis, 1998), and so competition for food is minimal.

In the absence of clouded leopards, the overlaps between species' activity patterns clearly differed (Fig. 5). The dyads including sambar occur mainly on the right of the *x*-axis where overlaps increased in the absence of clouded leopards. Increased contest competition between sambar deer and other ungulates (with the exception of the greater mouse deer) due to increased overlap in activity cycles, however, is unlikely for these very dispersed resources.

Temporal interactions are only one facet of the predator-prey relationship; spatial shifts of predator and prey populations relative to each other will play a role. The ecology of fear concept (Brown, Laundré & Gurung, 1999) codifies the naturalistic observation that prey species are likely to avoid areas

(even those good for foraging) where they are particularly vulnerable to predators. We have presented data consistent with the possibility of temporal avoidance of clouded leopards by bearded pigs, and perhaps by greater mouse deer. The next priority is to find out whether they, or indeed any of the species that did not exhibit a temporal shift, adapt their use of space to the risk of predation by clouded leopards.

Acknowledgements

We are indebted to our numerous field assistants and volunteers, Glen Reynolds and the Royal Society's SEARRP, Benoit Goossens and staff at Danau Girang Field Centre, for logistical support. We thank Norman Owen-Smith and an anonymous referee for helpful comments on earlier drafts. We thank our local collaborators Datin Maryati Mohammed, Professor Abdul Hamid, and Dr. Henry Bernard, Universiti Malaysia Sabah, and Soffian Abu Bakar, Sabah Wildlife Department. We thank Yayasan Sabah, Sabah Wildlife Department, Sabah Forestry Department, Danum Valley Management Committee, the FACE foundation, the State Secretary, the Sabah Chief Minister's Department, and the Prime Minister's Department (EPU) for research permission. This research was primarily funded by the Darwin Initiative, U.K. Department for Environment Food and Rural Affairs, Panthera, and a private donation from the Kaplan family to D.W.M. Additional funding was provided by the Clouded Leopard Project/Point Defiance Zoo and Aquarium, Felidae Conservation Fund, HGH Wills, International Trust for Nature Conservation, and Wild About Cats. During the preparation of this paper, J.R. was funded by grants to D.W.M. from the Robertson Foundation and the Wood-spring Trust.

References

- Barrette, C. (1977). Some aspects of behaviour of muntjacs in Wilpattu National Park. *Mammalia* **41**, 1–34.
- Bowyer, R.T., Van Ballenberghe, V., Kie, J.G. & Maier, J.A.K. (1999). Birth-site selection by Alaskan moose: maternal strategies for coping with a risky environment. *J. Mammal.* **80**, 1070–1083.
- Brown, J.S., Laundré, J.W. & Gurung, M. (1999). The ecology of fear: optimal foraging, game theory, and trophic interactions. *J. Mammal.* **80**, 385–399.
- de Bruyn, A.M.H. & Meeuwig, J.J. (2001). Detecting lunar cycles in marine ecology: periodic regression versus categorical ANOVA. *Mar. Ecol. Prog. Ser.* **214**, 307–310. [6]
- Carbone, C., Mace, G.M., Roberts, S.C. & Macdonald, D.W. (1999). Energetic constraints on the diet of terrestrial carnivores. *Nature* **402**, 442–442.
- Cheyne, S.M. & Macdonald, D.W. (2011). Wild felid diversity and activity patterns in Sabangau peat-swamp forest, Indonesian Borneo. *Oryx* **45**, 119–124.
- Christiansen, P. (2006). Sabertooth characteristics in the clouded leopard (*Neofelis nebulosa*, Griffiths 1821). *J. Morphol.* **267**, 1186–1198. [7]

- 1 Christiansen, P. & Adolfssen, J.S. (2005). Bite forces, canine
2 strengths and skull allometry in extant carnivores (Mam-
3 malia, Carnivora). *J. Zool.* **266**, 1–19.
- 4 Eccard, J., Pusenius, A., Sundell, J., Halle, S. & Ylönen, H.
5 (2008). Foraging patterns of voles at heterogeneous avian
6 and uniform mustelid predation risk. *Oecologia* **157**, 725–
7 734.
- 8 Fedriani, J.M., Palomares, F. & Delibes, M. (1999). Niche
9 relations among three sympatric Mediterranean carnivores.
10 *Oecologia* **121**, 138–148.
- 11 Fenn, M.G.P. & Macdonald, D.W. (1995). Use of middens by
12 red foxes: risk reverses rhythms of rats. *J. Mammal.* **76**,
13 130–136.
- 14 Gliwicz, J. & Drabowski, M.J. (2008). Ecological factors
15 affecting the diel activity of voles in a multi-species commu-
16 nity. *Ann. Zool. Fenn.* **45**, 242–247.
- 17 Gordon, C.H. & Stewart, A.E. (2007). The use of roads by
18 clouded leopards. *Cat News* **47**, 12–13.
- 19 Harmsen, B.J., Foster, R.J., Silver, S.C., Ostro, L.E.T. &
20 Doncaster, C.P. (2011). Jaguar and puma activity patterns
21 in relation to their main prey. *Mamm. Biol.* **76**, 320–324.
- 22 Jenny, D. & Zuberbühler, K. (2005). Hunting behaviour in
23 West African forest leopards. *Afr. J. Ecol.* **43**, 197–200.
- 24 Kamler, J.F., Johnson, A., Vongkhamheng, C. & Bousa, A.
25 (2012). The diet, prey selection and activity of dholes (*Cuon*
26 *alpinus*) in northern Laos. *J. Mammal.* **93**, 627–633.
- 27 Kawanishi, K. & Sunquist, M.E. (2004). Conservation status
28 of tigers in a primary rainforest of Peninsular Malaysia.
29 *Biol. Conserv.* **120**, 329–344.
- 30 Lemon, J. (2006). Plotrix: a package in the red light district of
31 R. *R-News* **6**, 8–12.
- 32 Linkie, M. & Ridout, M.S. (2011). Assessing tiger–prey inter-
33 actions in Sumatran rainforests. *J. Zool.* **284**, 224–229.
- 34 Macdonald, D.W., Mosser, A. & Gittleman, J.L. (2010). Felid
35 society. In *The biology and conservation of wild felids*: 125–
36 160. Macdonald, D.W. & Loveridge, A. (Eds). Oxford:
37 Oxford University Press.
- 38 Matsubayashi, H., Bosi, E. & Kohshima, S. (2003). Activity
39 and habitat use of lesser mouse-deer (*Tragulid javanicus*). *J.*
40 *Mammal.* **84**, 234–242.
- Matsuda, I., Tuuga, A. & Higashi, S. (2008). Clouded leopard
(*Neofelis diardi*) predation on proboscis monkeys (*Nasalis*
larvatus) in Sabah, Malaysia. *Primates* **49**, 227–231.
- Meek, P.D., Zewe, F. & Falzon, G. (2012). Temporal activity
patterns of the swamp rat (*Rattus lutreolus*) and other
rodents in north-eastern New South Wales, Australia. *Aust.*
Mammal. **34**, 223–233.
- Mills, M.G.L. & Shenk, T.M. (1992). Predator-prey relation-
ships: the impact of lion predation on wildebeest and zebra
populations. *J. Anim. Ecol.* **61**, 693–702.
- Mohamed, A., Samejima, H. & Wilting, A. (2009). Records of
five Bornean cat species from Deramakot Forest Reserve in
Sabah, Malaysia. *Cat News* **51**, 14–17.
- Owen-Smith, N. & Mills, M.G.L. (2008). Predator-prey size
relationships in an African large-mammal food web. *J.*
Anim. Ecol. **77**, 173–183.
- Payne, J. & Francis, C.M. (1998). *A field guide to the*
mammals of Borneo. Kota Kinabalu: The Sabah Society.
- Rabinowitz, A., Andau, P. & Chai, P.P.K. (1987). The
clouded leopard in Malaysian Borneo. *Oryx* **21**, 107–111.
- Ridout, M.S. & Linkie, M. (2009). Estimating overlap of daily
activity patterns from camera trap data. *J. Agric. Biol.*
Environ. Stat. **14**, 322–337.
- Schaller, G.B. (1967). *The deer and the tiger. A study of wild-*
life in India. Chicago and London: The University of
Chicago Press.
- Sunquist, M. & Sunquist, F. (2002). *Wild cats of the world*.
Chicago and London: The University of Chicago Press.
- Valeix, M., Loveridge, A.J., Chamaillé-Jammes, S., Davidson,
Z., Murindagomo, F., Fritz, H. & Macdonald, D.W.
(2009). Behavioural adjustments of African herbivores to
predation risk by lions: spatiotemporal variations influence
habitat use. *Ecology* **90**, 23–30.
- Van Schaik, C.P. & Griffiths, M. (1996). Activity periods of
Indonesian rain forest mammals. *Biotropica* **28**, 105–112.
- Yeager, C.P. (1991). Possible antipredator behaviour associ-
ated with river crossings by proboscis monkeys (*Nasalis*
larvatus). *Am. J. Primatol.* **24**, 61–66.
- Zar, J.H. (1996). *Biostatistics*. 3rd edn. ••, NJ: Prentice Hall,
Englewood Cliffs.

AUTHOR QUERY FORM

Dear Author,

During the preparation of your manuscript for publication, the questions listed below have arisen. Please attend to these matters and return this form with your proof.

Many thanks for your assistance.


Query References	Query	Remarks
1	AUTHOR: Please provide the current address.	
2	AUTHOR: Please check and confirm that the affiliation addresses are correct.	
3	AUTHOR: The (*) symbol has been replaced with (×) symbol in “significance of trigonometric × categoric terms.” Please check and confirm that this is correct.	
4	AUTHOR: de Bruyn <i>et al.</i> , 2003 has not been included in the Reference List, please supply full publication details.	
5	AUTHOR: Harmsen, 2011 has been changed to Harmsen <i>et al.</i> , 2011 so that this citation matches the Reference List. Please confirm that this is correct.	
6	AUTHOR: de Bruyn, Meeuwig, 2001 has not been cited in the text. Please indicate where it should be cited; or delete from the Reference List.	
7	AUTHOR: Christiansen, 2006 has not been cited in the text. Please indicate where it should be cited; or delete from the Reference List.	
8	AUTHOR: Christiansen, Adolfssen, 2005 has not been cited in the text. Please indicate where it should be cited; or delete from the Reference List.	
9	AUTHOR: Please supply the city location of publisher for Reference Zar 1996.	
10	AUTHOR: In Figure 4a and b, the label fonts are far too small, please increase their size.	
11	AUTHOR: The symbols * and ** in the sixth column of Table 2 have been changed to superscript (a) and (b), respectively, so that it does not confuse with the * symbols in the first column. Please check and confirm if this is OK, and provide the definitions each for (a) and (b) in Table 2 footnote.	

USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION


Required software to e-annotate PDFs: Adobe Acrobat Professional or Adobe Reader (version 8.0 or above). (Note that this document uses screenshots from Adobe Reader X)

The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/reader/>

Once you have Acrobat Reader open on your computer, click on the [Comment](#) tab at the right of the toolbar:



This will open up a panel down the right side of the document. The majority of tools you will use for annotating your proof will be in the [Annotations](#) section, pictured opposite. We've picked out some of these tools below:



1. Replace (Ins) Tool – for replacing text.




Strikes a line through text and opens up a text box where replacement text can be entered.

How to use it

- Highlight a word or sentence.
- Click on the [Replace \(Ins\)](#) icon in the Annotations section.
- Type the replacement text into the blue box that appears.

standard framework for the analysis of microeconomics. Nevertheless, it also led to the emergence of a new paradigm of strategic behaviour. The number of competitors in the industry is that the structure of the industry is a key component of the main components of the industry. At the level, are expected to be important works on the industry by Shiraz (M henceforth) we open the 'black b



2. Strikethrough (Del) Tool – for deleting text.



Strikes a red line through text that is to be deleted.

How to use it

- Highlight a word or sentence.
- Click on the [Strikethrough \(Del\)](#) icon in the Annotations section.

there is no room for extra profits and the number of competitors are zero and the number of (net) values are not determined by Blanchard and ~~Kiyotaki~~ (1987), perfect competition in general equilibrium. The effects of aggregate demand and supply in the classical framework assuming monopoly are an exogenous number of firms

3. Add note to text Tool – for highlighting a section to be changed to bold or italic.



Highlights text in yellow and opens up a text box where comments can be entered.

How to use it

- Highlight the relevant section of text.
- Click on the [Add note to text](#) icon in the Annotations section.
- Type instruction on what should be changed regarding the text into the yellow box that appears.

dynamic responses of mark ups consistent with the **VAR** evidence

sation y Ma and on n to a stent also with the demand-



4. Add sticky note Tool – for making notes at specific points in the text.




Marks a point in the proof where a comment needs to be highlighted.

How to use it

- Click on the [Add sticky note](#) icon in the Annotations section.
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the yellow box that appears.

and supply shocks. Most of the number of competitors and the impact is that the structure of the sector



USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION


5. Attach File Tool – for inserting large amounts of text or replacement figures.



Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it

- Click on the [Attach File](#) icon in the Annotations section.
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.



6. Add stamp Tool – for approving a proof if no corrections are required.




Inserts a selected stamp onto an appropriate place in the proof.

How to use it

- Click on the [Add stamp](#) icon in the Annotations section.
- Select the stamp you want to use. (The [Approved](#) stamp is usually available directly in the menu that appears).
- Click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

of the business cycle, starting with the
 on perfect competition, constant ret
 production. In this environment goods
 extra profits and the market for marke
 he market for goods is determined by the model. The New-Key
 otaki (1987), has introduced produc
 general equilibrium models with nomin
 and market-clearing. Most of this literat




7. Drawing Markups Tools – for drawing shapes, lines and freeform annotations on proofs and commenting on these marks.

Allows shapes, lines and freeform annotations to be drawn on proofs and for comment to be made on these marks..

How to use it

- Click on one of the shapes in the [Drawing Markups](#) section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, move the cursor over the shape until an arrowhead appears.
- Double click on the shape and type any text in the red box that appears.



For further information on how to annotate proofs, click on the [Help](#) menu to reveal a list of further options:

